

UNCLASSIFIED

AD **407 014**

DEFENSE DOCUMENTATION CENTER

FOR

SCIENTIFIC AND TECHNICAL INFORMATION

CAMERON STATION, ALEXANDRIA, VIRGINIA



UNCLASSIFIED

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

⑤ 70 900

⑦ 63-4-

407014

ALASKAN AIR COMMAND
ARCTIC AEROMEDICAL LABORATORY
FORT WAINWRIGHT

- ⑥
- ⑦ N
- ⑧ N
- ⑨ N
- ⑩ S
- ⑪ N
- ⑫ N
- ⑬
- ⑭
- ⑮
- ⑯
- ⑰
- ⑱
- ⑲

AD No. _____
DDC FILE COPY

MECHANISM OF ADAPTATION TO A THREONINE-DEFICIENT DIET,
PART IV

⑩ by Dorothy Arata,

TECHNICAL NOTE
⑩ AAL-TN-61-2

JAN 1 1963
RECEIVED
TISIA A

④ 4/10

407 014

⑪ OCTOBER 1961

(b) MECHANISM OF ADAPTATION TO A THREONINE-DEFICIENT DIET,

IV. Effect of age and weight on threonine deficiency symptoms in the rat, *all caps*

Dorothy Arata
Dept. of Foods and Nutrition
Michigan State University,

✓ When weanling rats are fed a threonine-deficient diet, fat deposits in the livers reach a maximum in approximately 2 weeks. This phase is followed by a period of recovery wherein the concentration of fat is slowly decreased. *tp*

The mechanisms controlling these responses have not been established. It is necessary to attempt to define the nature of the biochemical responses to a threonine deficiency.

This study was designed to measure whether or not the age and weight of the rat at the time of feeding the low protein, threonine-deficient diet exerted any influence on the severity of the resulting fatty liver syndrome.

EXPERIMENTAL

Eighty male weanling rats of the Sprague-Dawley strain were used in this experiment. The animals were divided into two major dietary groups. Rats in Group I were fed a 9% casein diet supplemented with 0.36% DL-threonine, while rats in Group II were fed the identical diet except that no threonine was added. The per cent composition of these diets has been presented in previous reports. In this experiment, however, an additional variable was introduced: a "pre-experimental" phase wherein the animals were fed Rockland pellets for varying periods of time. See Table I for clarification of the experimental design. Groups I₀ and II₀ served as control groups in this study of the effect of the age of the animal on the severity of the fatty liver syndrome. Food and water were allowed ad libitum.

After each group of rats had been fed the casein diet for a period of 2 weeks, the animals were stunned with a sharp blow on the head and decapitated. Livers were removed from the carcass, weighed, and homogenized. The homogenates were dried and ground, and samples weighing approximately 1 g were continuously extracted with ether for 3 hours.

Blood samples were taken from representative rats in each group. Electrophoretic patterns were run from serum prepared from each sample of whole blood.

RESULTS

Liver fat data are presented in Table II.

Livers from all rats fed casein diets deficient in threonine (~~group II~~) had significantly greater concentrations of fat than did the control animals. These differences were significant at the 1% level (~~$P < 0.01$~~) regardless of the length of the "pre-experimental feeding trial," (~~Table II~~). However, the absolute amount of fat deposited in livers from rats fed a threonine-deficient diet decreased as the length of the pellet-feeding phase increased. The same trend, though on a much smaller scale, was observed in the control rats (~~group I~~).

DISCUSSION

A quantity of liver fat significantly greater than in control animals was induced by feeding a threonine-deficient diet for 2 weeks to rats 21, 28, 35 and 42 days old. However, the amount of fat deposited varied directly with the age of the animal.

Since this experiment was designed as a pilot study, the data are insufficient to support the suggestion that the liver fat curve described in a threonine deficiency state is a function of age of the animal. Since the animals did not develop fatty livers to the same extent in all threonine-deficient groups, the age of the rat must play an important role in the development of this syndrome. But since complete data on the rate of fat deposition within a dietary treatment are not available for each group, it is not possible to assay the effect of a low protein, threonine-deficient diet on older animals. It is possible that in an older animal the peak in liver fat deposition induced by a threonine deficiency may occur more or less rapidly than in a younger rat.

A more extensive experiment is now in progress to attempt to explore these facets. The experimental groups defined in this paper have been used. However, each group will be studied separately through a long-term period (6 weeks). Rats will be studied with respect to liver fat deposition, growth, activity of serum transaminase, and concentration of serum proteins.

TABLE I
Experimental Design

Group *	Days on Pellets	Days on 9% Casein Diet	Threonine Supplement
I ₀	0	14	+
II ₀	0	14	--
I ₁	7	14	+
II ₁	7	14	--
I ₂	14	14	+
II ₂	14	14	--
I ₃	21	14	+
II ₃	21	14	--

* Each group was composed of 10 weanling rats.

TABLE II
Liver Fat Data for Rats Fed a Pellet Ration and
9% Casein With and Without Threonine Supplement

Weeks fed Pellets	% Dry Wt.	
	Group I (0.36% DL-Threonine)	Group II (No Threonine Supp.)
0	11.2 ± 0.6 *	21.0 ± 1.4 *
1	10.4 ± 0.5	18.4 ± 1.0
2	8.2 ± 0.6	15.4 ± 1.0
3	7.5 ± 0.3	10.6 ± 0.5

* Standard error of mean.